

Atrazine use (total lbs 1998-2002)

<u>Highest (82% of Total lbs)</u>	<u>Mid (17% of Total lbs)</u>	<u>Lowest (1% of Total lbs)</u>
FLORIDA	ALABAMA	ARIZONA
ILLINOIS	ARKANSAS	CALIFORNIA
INDIANA	COLORADO	CONNECTICUT
IOWA	DELAWARE	IDAHO
KANSAS	GEORGIA	MAINE
KENTUCKY	LOUISIANA	MASSACHUSETTS
MICHIGAN	MARYLAND	MONTANA
MINNESOTA	MISSISSIPPI	NEVADA
MISSOURI	NEW YORK	NEW HAMPSHIRE
NEBRASKA	NORTH CAROLINA	NEW JERSEY
OHIO	OKLAHOMA	NEW MEXICO
TEXAS	PENNSYLVANIA	NORTH DAKOTA
WISCONSIN	SOUTH CAROLINA	OREGON
	SOUTH DAKOTA	RHODE ISLAND
	TENNESSEE	UTAH
	VIRGINIA	VERMONT
		WASHINGTON
		WEST VIRGINIA
		WYOMING

<u>Birth Defect*</u>	
Spina	Hydro
Circul	Micro
Cleftlip	Nervous
Adactyly	Heart
Musculo	Rectal
Down	Omphalo
Tracheo	Genital
Gastro	Renalage
Urogen	Hernia
Clubfoot	Chromo
Anen	

*Abbreviation codes used by CDC

Birth Defects per 100,000 Births

